

Infertility Cycle

- Day 1: is always the first day of your menstrual cycle (the first day of bleeding). This is the day you will call to set up your appointments.
- Day 3: come in for labs to check hormone levels.
- Days 3-7: take your Clomid (Clomiphine Citrate) as directed.
- Days 3, 5, 7, and 9: inject 75 IU of Repronox - OR - Menopur intramuscularly or subcutaneously. (around \$200 cost per month)
- Days 7-10: schedule HSG if needed. (Do not forget to take Doxycycline every 12 hours the day before, day of, and day after the procedure).
- Day 10: come in for labs to check hormone levels.
- Days 8-12: take your Estradiol, 3 tablets, twice each day.
- Day 12: come in for your sonogram to check follicles.
 - Be sure to do your ovulation predictor test.
 - Bring your Novarel injection with you to this visit.
 - Have intercourse 24 hours after your injection every 12 hours for 48 hours -OR- come in 36 hours after injection for insemination.
- 3 days after ovulation until 14 days after ovulation: take your Progesterone, either oral pills or vaginal gel.
- Day 21 -OR- 7 days after ovulation: come in for labs to check Progesterone level.
- Day 28 -OR- 14 days after intercourse: come in for labs to check Pregnancy test if you have not started menses.